

Young Investigators Awards

CORONARY PHYSIOLOGICAL ADAPTATIONS DURING PERCUTANEOUS CORONARY INTERVENTION IN THE PRESENCE OF LEFT VENTRICULAR DYSFUNCTION: INSIGHTS FROM CORONARY WAVE INTENSITY ANALYSIS

Special Session
North, Room 120
Sunday, March 10, 2013, 11:00 a.m.-11:15 a.m.

Session Title: Young Investigator Awards Competition: Physiology, Pharmacology and Pathology
Abstract Category: Physiology, Pharmacology, Pathology
Presentation Number: 403-5

Authors: *Kalpa De Silva, Balrik Kailey, Jordi Alastruey-Armon, Matthew Lumley, Michael Marber, Sven Plein, Simon Redwood, Divaka Perera, St. Thomas' Hospital, King's College London, London, United Kingdom*

Introduction: Left ventricular (LV) dysfunction is often seen in acute coronary syndromes (ACS) and high-risk percutaneous coronary intervention (PCI). Wave intensity analysis (WIA) quantifies energy flux within the circulation. We used WIA to examine the hemodynamic perturbations following ACS and PCI with adjunctive intra-aortic balloon pump (IABP).

Methods: Patients with NSTEMI, or scheduled PCI with >2 vessel disease and LVEF <40% had simultaneous intra-coronary pressure-Doppler assessment. In the ACS group the target vessel (TV) and a remote reference vessel were assessed with LV function and late-gadolinium enhancement (LGE) assessed by cardiac MRI, pre- and 3-months post-PCI. In the PCI+IABP group, TV was assessed during unassisted and 40cc-IABP-assisted settings.

Results: 33 patients were recruited, 22 NSTEMI (LVEF $56 \pm 11\%$ and LGE mass $8.4 \pm 6.0\%$) and 11 PCI+IABP (LVEF $34 \pm 7\%$). TV BEW inversely correlated to LGE ($r = -0.81$, $p < 0.0001$) and predicted regional LV recovery ($r = 0.68$, $p = 0.001$). By ROC analysis, BEW predicted functional recovery with an accuracy of 88%. IABP resulted in a 12.7% ($p = 0.04$) and 28% ($p < 0.001$) increase in diastolic pressure, and flow velocity (DFV) respectively. A WIA derived early-diastolic wave, corresponding with IABP inflation, was proportional to DFV ($r = 0.72$, $p = 0.001$).

Conclusion: BEW correlates with magnitude and location of infarction, and may predict functional recovery following ACS. WIA uniquely assesses the physiological efficacy of mechanical assist devices.

